

Swift GRB GRB071010B : outlier of the $E_{\text{peak}}^{\text{src}} - E_{\gamma}$ and
 $E_{\text{iso}} - E_{\text{peak}}^{\text{src}} - t_{\text{jet}}^{\text{src}}$ correlations

Yuji URATA^{1,2,3}, Kuiyun HUANG³, Myungshin IM⁴ Induk LEE^{4,1} Jinsong DENG⁵,
 WingHuen IP¹, Hans KRIMM^{6,7}, Xin LIPING⁵, Masanori OHNO⁸, Yulei QIU⁵,
 Satoshi SUGITA^{9,10}, Makoto TASHIRO², Jianyan WEI⁵, Kazutaka YAMAOKA¹⁰, and
 Weikang ZHENG⁵

ABSTRACT

We present multi-band results for GRB071010B based on *Swift*, *Suzaku*, and ground-based optical observations. This burst is an ideal target to evaluate the robustness of the $E_{\text{peak}}^{\text{src}} - E_{\text{iso}}$ and $E_{\text{peak}}^{\text{src}} - E_{\gamma}$ relations, whose studies have been in stagnation due to the lack of the combined estimation of $E_{\text{peak}}^{\text{src}}$ and long term optical monitoring. The joint prompt spectral fitting using *Swift*/Burst Alert Telescope and *Suzaku*/Wide-band All sky Monitor data yielded the spectral peak energy as $E_{\text{peak}}^{\text{src}}$ of $86.5_{-6.3}^{+6.4}$ keV and E_{iso} of $2.25_{-0.16}^{+0.19} \times 10^{52}$ erg with $z = 0.947$. The optical afterglow light curve is well fitted by a simple power law with

¹Institute of Astronomy, National Central University, Chung-Li 32054, Taiwan; urata@astro.ncu.edu.tw

²Department of Physics, Saitama University, Shimo-Okubo, Saitama, 338-8570, Japan

³Academia Sinica Institute of Astronomy and Astrophysics, Taipei 106, Taiwan

⁴Center for the Exploration of the Origin of the Universe, Department of Physics & Astronomy, FPRD, Seoul National University, Shillim-dong, San 56-1, Kwanak-gu, Seoul, Korea

⁵National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China

⁶NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

⁷Universities Space Research Association, 10211 Wincopin Circle, Suite 500, Columbia, MD 21044, USA

⁸Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Sagami-hara, Kanagawa 229-8510, Japan

⁹The Institute of Physical and Chemical Research (RIKEN), 2-1 Hirosawa, Wako, Saitama, 351-0198, Japan

¹⁰Department of Physics and Mathematics, Aoyama Gakuin University, 5-10-1, Fuchinobe, Sayamihara 229-8558, Japan

temporal index $\alpha = -0.60 \pm 0.02$. The lower limit of temporal break in the optical light curve is 9.8 days. Our multi-wavelength analysis reveals that GRB071010B follows $E_{\text{peak}}^{\text{src}} - E_{\text{iso}}$ but violates the $E_{\text{peak}}^{\text{src}} - E_{\gamma}$ and $E_{\text{iso}} - E_{\text{peak}}^{\text{src}} - t_{\text{jet}}^{\text{src}}$ at more than the 3σ level.

Subject headings: gamma rays: bursts — gamma rays: observation

1. Introduction

A few tight correlations linking several properties of gamma-ray bursts (GRBs), namely the spectral peak energy, the total radiated energy, and the afterglow break time, have been discovered with pre-*Swift* GRBs. Amati et al. (2002) found the empirical relation between the rest-frame spectral peak energy of prompt emission $E_{\text{peak}}^{\text{src}}$ (here, we use the $E_{\text{peak}}^{\text{src}}$, $E_{\text{peak}}^{\text{obs}}$ for the values in the rest and observer frames, respectively) and the isotropic-equivalent energy released during the prompt phase E_{iso} . Ghirlanda et al. (2004a) showed a tight empirical relation between the $E_{\text{peak}}^{\text{src}}$ and the collimation-corrected energy E_{γ} based on a jet interpretation of the temporal break in the optical afterglow light curve. Liang & Zhang (2005) found the three-dimensional correlation in the $E_{\text{iso}} - E_{\text{peak}}^{\text{src}} - t_{\text{jet}}^{\text{src}}$ plane. The existence of these correlations could uncover crucial properties of GRB physics which are not yet fully understood. Recently, there have been studies of outliers to these relations (e.g., Campana et al 2007, Sato et al. 2007, Ghirlanda et al. 2007). Using *Swift*/XRT observations, Sato et al. (2007) showed that there are cases with continuous X-ray afterglow light curves where no jet break is seen at the expected break time derived from the $E_{\text{peak}}^{\text{src}} - E_{\gamma}$ relation. However, some X-ray afterglow light curves may not show jet breaks together with the optical afterglow at the expected time, because there is a possibility that X-ray and optical emission come from different regions and mechanisms even in the normal decay phase (e.g., Urata et al. 2007a, Liang et al 2007, Huang et al. 2007, Ghisellini et al. 2009). Therefore, it is critical to check the aforementioned relations by combining prompt γ -ray wide-band spectroscopy and optical long-term monitoring.

In this Letter, we present joint spectral analysis of the prompt gamma-ray and systematic optical follow-up results for GRB 071010B. This event was detected and localized by *Swift* (Markwardt et al. 2007a). The *Suzaku*/Wide-band All sky Monitor (WAM; Yamaoka et al. 2009) and Konus/*WIND* (Golenetskii et al. 2007) also detected this main burst. Thanks to the wide energy band of the *Suzaku*/WAM (Yamaoka et al. 2009), the joint spectral fittings between the *Swift*/Burst Alert Telescope (BAT) and the *Suzaku*/WAM data yield better constraint for the determination of E_{peak} . The optical afterglow was discovered by Oksanen (2007) and observed by many telescopes from the early stage (e.g., Wang et al.

2007). The redshift was determined to be $z = 0.947$ by Cenko et al. (2007) using the Gemini North telescope. In the frame work of EAFON (Urata et al. 2003, 2005), we have performed long term monitoring to determine the jet break time. Therefore, GRB071010B is one of the most suitable event to evaluate $E_{\text{peak}}^{\text{src}} - E_{\text{iso}}$, $E_{\text{peak}}^{\text{src}} - E_{\gamma}$ and $E_{\text{iso}} - E_{\text{peak}}^{\text{src}} - t_{\text{jet}}^{\text{src}}$ in *Swift* era. All the errors are quoted at the 90% statistical confidence level in this paper.

2. Observations

2.1. *Swift*/BAT

The *Swift* /BAT triggered and located GRB 071010B (trigger ID=293795) at 20:45:47 (T_0) UT on 2007 October 10. The on-board localization was distributed 13 sec after the trigger and its position was reported as RA= $10^{\text{h}}02^{\text{m}}07.5$, Dec= $+45^{\circ}44'0''$, with an uncertainty of $3'$. The spacecraft did not slew promptly to the burst position because automated slewing was disabled due to ongoing recovery from a gyro anomaly (Markwardt et al. 2007a). After 6800 s of the trigger, *Swift*/XRT started observing the field and detected a bright X-ray counterpart at R.A.= $10^{\text{h}}02^{\text{m}}09.2$, decl.= $+45^{\circ}43'52''.2$, with an uncertainty of $5''$. Markwardt et al. (2007b) reported the mask-weighted light curve which shows a pre-trigger pulse starting at $T_0 \sim -45$ s, peaking at $T_0 \sim -20$ s, and returning almost to background at $T_0 \sim -8$ s. The main FRED pulse started at $T_0 \sim 2$ s and ended around $T_0 \sim 60$ s. A small third soft peak started at $T_0 + 95$ s and was terminated by the planned spacecraft slew (Markwardt et al. 2007b). The two peaks at $T_0 - 45$ s and $T_0 + 95$ s are much weaker than the main peak, thus they are not visible in Figure 1.

2.2. *Suzaku*/WAM

The *Suzaku*/WAM was triggered at 20:45:49 ($T_0 + 2$ s) UT on 2007 October 10. The *Suzaku*/WAM is the active shield of the Hard X-ray detector (Takahashi et al. 2007; Kokubun et al. 2007) onboard the 5th Japanese X-ray satellite *Suzaku* (Mitsuda et al. 2007). It consists of large area, thick BGO crystals and is also designed to monitor the all sky from 50 keV to 5 MeV by a large effective area. The large effective area from 300 keV to 5 MeV (400cm^2) surpasses those of other currently operating experiments with spectral capability, and enables us to perform wide-band spectroscopy of GRBs with high sensitivity.

As shown in Figure 1, the WAM light curve also shows a single FRED-like peak starting at $T_0 - 1$ s, ending at $T_0 + 8$ sec. The duration of the WAM signal is $T_{90} = 5$ s. This spike was also detected by both the *Swift*/BAT, and the Konus/*WIND*. There is no signif-

icant signal from the pre-cursor and the weak soft tail seen in the *Swift*/BAT light curve (Markwardt et al. 2007b) during the time coverage.

2.3. Optical Follow-ups

We carried out follow-up observations of the GRB071010B optical afterglow at the Mt. Lemmon, Arizona, USA and the Xinglong, China observatories within the framework of the EAFON (Urata et al. 2003, 2005). Using the robotic 1 m telescope and a $2k \times 2k$ CCD camera at the Mt. Lemmon observatory operated by the Korea Astronomy Space Science Institute (I. Lee et al. in preparation,; Han et al. 2005), we have made *B*-, *V*-, and *R*-band imaging observations with 300 s exposures, starting at 10:51:14.6 UT on 2007 October 11 (0.5871 days after the burst). In all the bands, we detected the afterglow clearly. We have also monitored the afterglow on October 12, 16, 19, 21, 25, and 26 in the *R*-band. Additional *V*-band observations were performed using the EST 1.0 m telescope at the Xinglong Observatory (Zheng et al. 2008), starting at 2007 October 11 (19:08 UT; 0.971 days after the burst). The field of view is $11'.4 \times 11'.1$, and the pixel size is $0''.51$ square.

The *i'*- and *z'*-band observations were acquired with the 3.6-m CFHT using MegaCam on 2008 October 26, and November 6 (approximately one year after the burst), respectively. These deeper images show the host galaxy with $i' = 23.63 \pm 0.14$ AB mag, $z' = 22.73 \pm 0.11$ AB mag at R.A. = $10^h02^m09.274$, decl. = $+45^\circ43'49''.69$.

3. Analysis and Results

3.1. *Swift*/BAT Prompt Emission

The BAT data were analyzed using the standard BAT analysis software distributed within HEADAS v6.4. Using batgrbproduct v2.41, the mask-weighted BAT light curves were created in the standard four energy bands, 15 – 25, 25 – 50, 50 – 100, 100 – 350 keV, and the duration was determined as $T_{90} = 36.0 \pm 2.4$ s. Figure 1 shows 15 – 50 keV and 50 – 150 keV light curves. Response matrices were generated with the task batdrngen using the latest spectral redistribution matrices. The time-averaged spectrum (15 – 150 keV) from $T_0 - 35.7$ to $T_0 + 24.2$ s is well fitted by a power law with an exponential cutoff. This fit yields a photon index of 1.53 ± 0.22 , and E_{peak}^{obs} of 52.0 ± 6.4 keV. A fit to a simple power law gives a photon index of 2.01 ± 0.05 ($\chi^2/\nu = 0.82$ for $\nu = 57$).

3.2. *Suzaku*/WAM Prompt Emission

The WAM spectral and temporal data were extracted using `hxdmkwamlc` and `hxdmkwamspec` in the HEADAS version 6.4. The background was estimated using the fitting model described in Sugita et al. (2009). Response matrices were generated by the WAM response generator as described in Ohno et al. (2008). The time-averaged spectrum (150 – 1000 keV) from $T_0 - 0.78$ to $T_0 + 24.2$ s was well fitted by a single power law with a photon index of $2.64^{+0.26}_{-0.22}$ ($\chi^2/\nu = 1.07$ for $\nu = 10$). This result is consistent with the *Swift*/BAT power law with an exponential cutoff fitting with the photon index 1.53 ± 0.22 , and E_{peak}^{obs} of 52.0 ± 6.4 keV.

3.3. BAT and WAM Joint Analysis

In order to better constrain the spectral peak energy, we perform joint fitting between *Swift*/BAT and *Suzaku*/WAM. Figure 1 shows the overlapping time regions (from $T_0 - 0.780$ to $T_0 + 24.2$ s) for this fitting. As shown in Figure 2, the spectrum is well fitted with the Band function (Band et al 1993). The fitting yields a low-energy photon index of $1.19^{+0.29}_{-0.23}$, a high-energy photon index of $2.30^{+0.06}_{-0.07}$ and E_{peak}^{src} of $86.5^{+6.4}_{-6.3}$ keV ($\chi^2/\nu = 0.59$ for $\nu = 87$). This result is consistent with that of *Konus/WIND* (Golenetskii et al. 2007). We have also estimated the E_{iso} as $2.25^{+0.19}_{-0.16} \times 10^{52}$ erg, assuming cosmological parameters: $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.27$, and $\Omega_\Lambda = 0.73$.

3.4. Optical Afterglow

A standard routine including bias subtraction and flat-fielding corrections was employed to process the data using the *IRAF* package. The DAOPHOT package was used to perform aperture photometry of the GRB images. For the photometric calibration of the afterglow, several calibration stars reported by Henden (2007) were chosen. In Figure 3, we plot the *Rc*-band light curve of the GRB071010B afterglow based on our photometry. The data contain our LOAO measurements, TAOS observations (Wang et al. 2008) with the re-calibrated GCN measurements (Oksanen (2007); Kann et al. (2007a,b,c); Kocevski et al. (2007); Klunko et al. (2007)). The light curve between 0.02 and 10.6 day is well fitted by a single power law with index $\alpha = -0.60 \pm 0.02$ ($\chi^2/\nu = 0.89$ for $\nu = 37$), defined by $F(t) \propto t^\alpha$, where $F(t)$ is the flux in the *Rc*-band at time t after the BAT trigger time T_0 . The single power-law index is consistent with that of the TAOS measurement ($\alpha \sim -0.51$) reported by Wang et al. (2008). This fitting also excludes the possible 3.4 days temporal break reported

by Kann et al (2007) and Im et al (2007) based on GCN data point measurements. We have also confirmed that the host-galaxy brightness ($i' = 23.63$ mag, $z' = 22.73$ mag) derived by the CFHT observations is insufficient to affect this fitting. We consider both red and typical host-galaxy colors to evaluate R -band brightness. Although the late-time light curve could be fitted by a single power-law, it is interesting to note that a possible temporal break occurred at around 9 days after the burst. We tried to fit the 0.02–10.6 days light curve with a broken power law function expressed as

$$F_\nu(t) = \frac{F_\nu^*}{[(t/t_b)^{\alpha_1} + (t/t_b)^{\alpha_2}]}, \quad (1)$$

where α_1 and α_2 are the power-law indices before and after the break, F_ν^* is the flux at the break, and t_b is the break time. The fitting yields $\alpha_1 = -0.54 \pm 0.03$, $\alpha_2 = -3.03 \pm 1.15$ and $t_b = 9.81 \pm 1.00$ ($\chi^2/\nu = 0.52$ for $\nu = 35$). When we fix α_2 as -2, the fitting provides the break time of $t_b = 9.61 \pm 1.49$. Therefore, these results imply that the temporal break should be around or later than 9.8 days after the burst.

4. Discussion

Spectral parameters of the prompt emission of GRB071010B are well constrained by our joint fitting of *Swift*/BAT and *Suzaku*WAM. The measured values of current event ($E_{peak}^{src} = 86.5_{-6.3}^{+6.4}$ keV and $E_{iso} = 2.25_{-0.16}^{+0.19} \times 10^{52}$ erg) well follow the $E_{peak}^{src} - E_{iso}$ relation. Furthermore, our systematic long term (10 days) optical monitoring observation suggests a possible jet break at later than ~ 9.8 days after the burst. The wealth of the multi-wavelength data with a good temporal coverage makes GRB071010B one of the best targets to evaluate the $E_{peak}^{src} - E_\gamma$ and the $E_{iso} - E_{peak}^{src} - t_{jet}^{src}$ relations which studies have been in stagnation due to the lack of the E_{peak}^{src} estimation and long term optical monitoring.

In order to evaluate the $E_{peak}^{src} - E_\gamma$ relation for the optical light curve, we invert this relation to predict the jet break time as described in Sato et al. (2007). The expected jet break time is expressed as

$$t_{jet} = 389 (1+z) \left(\frac{n}{3 \text{ cm}^{-3}} \right)^{-\frac{1}{3}} \left(\frac{\eta_\gamma}{0.2} \right)^{-\frac{1}{3}} E_{iso,52}^{-1} \left(\frac{E_{peak}^{src}}{A} \right)^{1.89} \text{ day}, \quad (2)$$

where n and η_γ are the number density of the ambient medium and the efficiency of the shock, respectively. Here, we use the Ghirlanda relation as $E_{peak}^{src} = A E_{\gamma,52}^{0.70}$, allowing A to vary within the 3σ level. The value of n is in general within $1 < n < 30 \text{ cm}^{-2}$ (Panaiteescu & Kumar 2001, 2002). Following the assumption made by Ghirlanda et al. (2004a) for most of their sample GRBs, we initially assume $n = 3 \text{ cm}^{-2}$ and $\eta_\gamma = 0.2$. As shown in Figure 3, the

optical light curve is well described with a single power law during the expected jet break time. We also tried to fit the optical light curve by fixing the jet break time as $t_j = 1.25$ days expected from Equation (2). The fitting yields $\alpha_1 = -0.30 \pm 0.06$ and $\alpha_2 = -0.91 \pm 0.04$ ($\chi^2/\nu=0.72$ for $\nu = 36$). Although this fitting is acceptable in terms of the χ^2 -fitting, an F -test indicates that this broken power-law model with $t_j = 1.25$ days over the single power law is not significant at the 90% confidence level. These temporal decay indices are inconsistent with those of typical jet break events. The temporal ($\alpha_x = 0.67$) and spectral ($\beta_x = 1.10$) relation of X-ray afterglow also support the sphere phase ($\alpha = 3/2\beta - 1 \sim 0.65$). These X-ray values come from UK's X-ray afterglow repository (Evans et al 2009). Thus, these results imply that GRB071010B is an outlier of the tight $E_{\text{peak}}^{\text{src}} - E_\gamma$ relation.

In Figure 4, we also plotted the $E_{\text{peak}}^{\text{src}} - E_\gamma$ relation together with GRB041006, GRB050904 and others reported by Ghirlanda et al. (2007). According to the detailed optical analysis of the GRB041006 afterglow (Urata et al. 2007b), this pre-*Swift* era event shows a chromatic optical plateau phase around $0.1 \sim 0.2$ days after the burst. Since Ghirlanda et al. (2007) regarded this chromatic plateau in the GRB041006 optical light curve as the jet break, we updated E_γ for this event. As shown in Figure 4, GRB071010B is a clear outlier of the tight correlation with more than 3σ deviation. GRB041006 is also possible outlier at the 2σ level. We have also tested the wind case. The estimated $E_{\gamma,w} > 2.8 \times 10^{50}$ erg from the jet break limit is inconsistent with Ghirlanda et al (2007) by over 3σ level. According to Sugita et al. (2009), GRB050904 is also an outlier of the $E_{\text{peak}}^{\text{src}} - E_\gamma$ relation, but it follows the $E_{\text{peak}}^{\text{src}} - E_{\text{iso}}$ relation.

These three events including a high-redshift GRB050904 imply the $E_{\text{peak}}^{\text{src}} - E_\gamma$ relation could have larger scatter than originally suggested. Since the $E_{\text{peak}}^{\text{src}} - E_\gamma$ correlation requires a small scatter to be used to standardize GRBs in a way similar to Type Ia supernovae (Ghirlanda et al. 2004b), these three events pose a problem to the cosmological application of the $E_{\text{peak}}^{\text{src}} - E_\gamma$ relation. Another implication is that these outliers have the relevant role of the circumburst environments for different origins (e.g., long and short bursts). In the GRB050904 case, the gap can be explained by a higher circumburst density which implies a massive stars origin. The required density is consistent with the estimations by optical and radio observations (Sugita et al. 2009). However, GRB071010B requires a 2 or 3 order smaller circumburst density ($n \sim 5.9 \times 10^{-3} \text{cm}^{-3}$) to be consistent with the tight correlation. This lower density environment implies the origin of GRB071010B is of short GRB, where typical circumburst density is $n = 10^{-2} \text{cm}^{-3}$ (e.g., Nakar 2007), even though the prompt duration of GRB071010B is longer than 2 s ($T_{90} = 36.0 \pm 2.3$ s).

We also tested the empirical $E_{\text{iso}} - E_{\text{peak}}^{\text{src}} - t_{\text{jet}}^{\text{src}}$ relation (Liang & Zhang 2005) expressed

as

$$E_{\text{iso}}/10^{52} = (0.85 \pm 0.21) \left(\frac{E_{\text{peak}}^{\text{src}}}{100\text{keV}} \right)^{1.94 \pm 0.17} \left(\frac{t_{\text{jet}}^{\text{src}}}{1\text{day}} \right)^{-1.24 \pm 0.23}, \quad (3)$$

where $t_{\text{jet}}^{\text{src}}$ is the jet break time in the rest frame. As shown in Figure 5, this relation yields the expected upper limit of isotropic total energy as $E_{\text{iso}} < 0.11 \times 10^{52}\text{erg}$ with the jet break time $t_{\text{jet}} > 9.8$ day. Therefore, GRB071010B is also an outlier of this model-independent relation at more than the 3σ level.

Since the tight correlations $E_{\text{peak}}^{\text{src}} - E_{\gamma}$ and $E_{\text{iso}} - E_{\text{peak}}^{\text{src}} - t_{\text{jet}}^{\text{src}}$ might indicate a crucial property of GRB physics (e.g., Thompson, (2006) and Thompson, Meszaros & Rees, (2007) for the $E_{\text{peak}}^{\text{src}} - E_{\gamma}$ relation), GRB071010B might be classified as a new category. However, there is also the possibility that these relations simply have larger dispersion than previous studies. Therefore, further study is required with large volume of good samples which can be built up by the joint analysis of *Swift*/BAT, *Suzaku*/WAM, *Konus*/*WIND*, *Fermi*/GBM and future missions such as the Space Variable Objects Monitor (SVOM) (Basa et al. 2008) with ground-based optical follow-ups.

We thank Bing Zhang for useful comments and discussions. M.I. and I.L. acknowledge the support by the Creative Research Initiatives grant R16-2008-015-01000-0 of MOST/KOSEF. This work is partly supported by grants NSC 98-2112-M-008-003-MY3 (Y.U.), NSC 96WFA0700264 (W.H.I.), Ministry of Education under the Aim for Top University Program NCU (W.H.I.), National Natural Science Foundation of China (No. 10673014), and National Basic Research Program of China (No. 2009CB824800). Access to the CFHT was made possible by the Ministry of Education and the National Science Council of Taiwan as part of the Cosmology and Particle Astrophysics (CosPA) initiative.

REFERENCES

- Amati, L., et al. 2002, *A&A*, 390, 81
- Basa, S., et al. 2008, in proceedings of the SF2A conference, Paris, 2008 (arXiv:0811.1154)
- Cenko, S. B., Cucchiara, A., Fox, D. B., Berger, E., & Price, P. A. 2007, *GRB Coordinates Network*, 6888, 1
- Campana, S., Guidorzi, C., Tagliaferri, G., Chincarini, G., Moretti, A., Rizzuto, D., & Romano, P. 2007, *A&A*, 472, 395
- Ghirlanda, G., Ghisellini, G., & Lazzati, D. 2004a, *ApJ*, 616, 331

- Ghirlanda, G., Ghisellini, G., Lazzati, D., & Firmani, C. 2004b, *ApJ*, 613, L13
- Ghirlanda, G., Nava, L., Ghisellini, G., and Firmani, G., 2007, *Å*, 466, 127
- Ghisellini, G., et al., 2009, *MNRAS* 393, 253
- Golenetskii et al. 2007, *GCN Circ.*, (2007) 6879
- Han, W., et al. 2005, *PASJ*, 57, 821
- Huang, K.Y., et al. 2007, *ApJ*, 654, L25
- Im, M., I. Lee, & Urata, Y. 2007, *GRB Coordinates Network*, 6897, 1
- Kann, D. A., Hoegner, C., & Filgas, R. 2007, *GRB Coordinates Network*, 6918, 1
- Kann, D. A., Laux, U., & Filgas, R. 2007, *GRB Coordinates Network*, 6923, 1
- Kann, D. A., et al. 2007, *GRB Coordinates Network*, 6935, 1
- Klunko, E., Marchenkov, A., Eselevich, M., Shulga, A., Volnova, A., & Pozanenko, A. 2007, *GRB Coordinates Network*, 6945, 1
- Kira, C., et al. 2007, *GRB Coordinates Network*, 6931, 1
- Kocevski, D., Perley, D. A., Bloom, J. S., Modjaz, M., & Poznanski, D. 2007, *GRB Coordinates Network*, 6919, 1
- Kokubun, M., et al. 2007, *PASJ*, 59, 53
- Liang, E. W., Zhang, B. B, Zhang, B, 2007 *ApJ*, 670, 565
- Liang, E. W., and Zhang, B, 2005 *ApJ*, 633, 611
- Markwardt, C., 2007, *GCN Circ.*, (2007) 6871
- Markwardt, C., Kennea, J., Mangano, V., Barthelmy, S. D., Burrows, D. N., Roming, P., & Gehrels, N. 2007, *GCNR*, 92, 1 (2007), 92, 1
- Mitsuda, K., et al. 2007, *PASJ*, 59, 1
- Nakar, E. 2007, *Phys. Rep.*, 442, 166
- Oksanen, A. 2007, *GRB Coordinates Network*, 6873, 1
- Ohno, M., et al. 2008, *PASJ*, 60, 361

- Sato, G., et al. 2007, ApJ, 657, 359
- Shirasaki, Y., et al. 2008, PASJ, 60, 919
- Sugita, S., et al. 2009, PASJ, in press
- Takahashi, T., et al. 2007, PASJ, 59, 35
- Thompson, C., 2006, ApJ, 651, 333
- Thompson, C., Meszaros, P., & Rees, M.J., 2007, ApJ, 666, 1012
- Urata, Y., et al. 2003, ApJ, 595, L21
- Urata, Y., et al. 2005, Nuovo Cimento C Geophysics Space Physics C, 28, 775
- Urata, Y., et al. 2007a, ApJ, 668, L95
- Urata, Y., et al. 2007b, ApJ, 655, L81
- Yamaoka, K., et al. 2009, PASJ, 61, S35
- Wang, J. H., Schwamb, M. & Huang K. Y. et al. 2008 ApJ, 679, L5
- Zheng, W. K., et al. 2008 ChJAA, 8, 693




Fig. 1.— Prompt X-ray and gamma-ray light curve of GRB071010B observed by *Swift*/BAT and *Suzaku*/WAM. The data in the time region between the dashed lines are used for the joint spectral analysis.




Fig. 2.— Broad band spectrum of GRB071010B prompt emission made by the joint analysis of the *Swift*/BAT and *Suzaku*/WAM data.




Fig. 3.— Rc -band afterglow light curve. The time region enclosed by the dashed (3σ) and the dash-dotted (5σ) lines indicates the expected jet break time from the $E_{\text{peak}}^{\text{src}} - E_{\gamma}$ relation.




Fig. 4.— $E_{peak}^{src} - E_\gamma$ relation including the data for GRB071010B and GRB041006 corrected for a homogeneous circumburst medium. The solid line shows the best-fit correlation derived by Ghirlanda et al. (2007). The dashed and dash-dotted lines indicate the 1σ and 3σ scatter of the correlation, respectively.




Fig. 5.— Plot of $\log \hat{E}_{\text{iso}}$ calculated by the empirical $E_{\text{iso}} - E_{\text{peak}}^{\text{src}} - t_{\text{jet}}^{\text{src}}$ relation (Liang & Zhang 2005) as compared with $\log E_{\text{iso}}$ derived from the observed fluence. The arrow indicates the upper limit of GRB071010B.